

Triplet-Phase Measurements in Crystallography - an Application of Triple Correlation Theory from Speckle Imaging


Qun Shen

Cornell High Energy Synchrotron Source (CHESS) and Department of Materials
Science and Engineering

Wilson Laboratory, Cornell University, Ithaca, New York 14853

E-mail: qs11@cornell.edu

We present a recent advance towards solving the phase problem in x-ray crystallography using a *reference-beam diffraction* (RBD) technique [1,2]. This technique incorporates the principle of three-beam interference into the popular oscillating-crystal method in crystallography and allows parallel recordings of many three-beam interference profiles on an area detector. From these measured interference profiles a large number of triplet-phases of the reflections involved can be obtained. These measured phases can then be used in conjunction with the direct methods to solve a crystal structure, without the need for heavy atoms.



The RBD phasing technique in fact can be viewed as a direct application of the triple correlation or bispectrum processing method in speckle imaging [3,4]. A recursive phase retrieval scheme can thus be applied on the triplet-phases measured in a single RBD data set to obtain potentially all individual structure-factor phases of a given structure. This is being investigated as an alternative strategy to solve the phase problem in x-ray crystallography.

This work is supported by NSF Grant DMR-9713424 through CHESS.

1. Q. Shen, *Phys. Rev. Lett.* 80, 3268 (1998); *Phys. Rev.* B59, 11109 (1999).
2. Q. Shen, S. Kycia, and I. Dobrianov, *Acta Cryst.* A56, 268 (2000).
3. A.W. Lohmann, G. Weigelt, and B. Wirnitzer, *Appl. Opt.* 22, 4028 (1983).
4. R.P. Millane, *J. Opt. Soc. Am.* A7, 394 (1990).